

Study of $B_c \rightarrow J/\psi K$ decays in the perturbative QCD approach

Xian-Qiao Yu*, Xiu-Li Zhou

School of Physical Science and Technology, Southwest University, Chongqing 400715, China

In this note, we calculate the branching ratio of $B_c \rightarrow J/\psi K$ in the framework of perturbative QCD approach based on k_T factorization. This decay can occur only via tree level diagrams in the Standard Model. We find that the branching ratio of $B_c \rightarrow J/\psi K$ is about $(1 - 3) \times 10^{-4}$. The large branching ratio and the clear signals of the final states make the measurement of $B_c \rightarrow J/\psi K$ easily at LHC-b experiments.

PACS numbers: 13.25.Hw, 12.38.Bx, 12.39.St

The B meson rare decays provide a good place for testing the Standard Model (SM), studying CP violation and looking for possible new physics beyond the SM. The theoretical studies of $B_{u,d}$ mesons decays have been studied widely in the literature, which are strongly supported by the detectors at the e^+e^- colliders, such as the CLEO, BaBar and Belle. At the era of the Large Hadron Collider(LHC), there is still a room for B physics. LHC beauty experiments(LHCb) will extend the B-physics results from the B factories by investigating decays of heavier B hadrons, such as B_s and B_c mesons. It is estimated that about 5×10^{10} B_c mesons can be produced per year at LHC[1, 2], so the studies of B_c meson rare decays are necessary in the next a few years, it will highlight the advantages of B physics.

In this paper, we study the rare decays $B_c \rightarrow J/\psi K$ in the Perturbative QCD approach (PQCD) [3]. In SM, $B_c \rightarrow J/\psi K$ decays occur through only the tree level diagrams and so there is no CP violation in this channel. Therefore, the measurements of this channel might provide a ground for investigating new physics effects.

For the decay $B_c \rightarrow J/\psi K$, the related effective Hamiltonian is given by [4]

$$H_{\text{eff}} = \frac{G_F}{\sqrt{2}} V_{us} V_{cb}^* \{C_1(\mu) O_1(\mu) + C_2(\mu) O_2(\mu)\} + H.c., \quad (1)$$

where $C_i(\mu)$ are Wilson coefficients at the renormalization scale μ and O_i are the local four-quark operators

$$O_1 = (\bar{b}_i c_j)_{V-A} (\bar{u}_j s_i)_{V-A}, \quad O_2 = (\bar{b}_i c_i)_{V-A} (\bar{u}_j s_j)_{V-A}. \quad (2)$$

Here i and j are $SU(3)$ color indices. Then the calculation of decay amplitude is to evaluate the hadronic matrix elements of the local operators.

In the PQCD approach, the decay amplitude can be written as:

$$\text{Amplitude} \sim \int d^4 k_1 d^4 k_2 d^4 k_3 \text{Tr}[C(t) \Phi_{B_c}(k_1) \Phi_\psi(k_2) \Phi_K(k_3) H(k_1, k_2, k_3, t)] e^{-S(t)}. \quad (3)$$

In our following calculations, the Wilson coefficient $C(t)$, Sudakov factor $S_i(t)$ ($i = B_c, J/\psi, K$) and the non-perturbative but universal wave function Φ_i can be found in the Refs. [5–10]. The hard part H are channel dependent but fortunately perturbative calculable, which will be shown below.

Fig. 1 shows the lowest order Feynman diagrams to be calculated in the PQCD approach where (a) and (b) are factorizable topology, (c) and (d) are nonfactorizable topology. After a straightforward calculation using the PQCD formalism Eq.(3), we obtain the sum contributions of (a) and (b)

$$\begin{aligned} M_a[C] = & \frac{8f_{B_c}}{\sqrt{2N_c}} \pi C_F M_{B_c}^2 \int_0^1 dx_2 \int_0^\infty b_1 db_1 b_2 db_2 \times \{[r_b r(1-r^2) \phi_\psi^t(x_2, b_2) - 2r(1-r^2) x_2 \phi_\psi^t(x_2, b_2) \\ & + (1-r^2) x_2 \phi_\psi^L(x_2, b_2) - 2r_b(1-r^2) \phi_\psi^L(x_2, b_2)] \alpha_s(t_a^1) h_a^{(1)}(x_2, b_1, b_2) \exp[-S_{B_c}(t_a^1) - S_{J/\psi}(t_a^1)] C(t_a^1) \\ & + [r^2(1-r^2)(r_c - 1) \phi_\psi^L(x_2, b_2) - r_c(1-r^2) \phi_\psi^L(x_2, b_2)] \alpha_s(t_a^2) h_a^{(2)}(x_2, b_1, b_2) \\ & \times \exp[-S_{B_c}(t_a^2) - S_{J/\psi}(t_a^2)] C(t_a^2)\}, \end{aligned} \quad (4)$$

* yuxq@swu.edu.cn




FIG. 1: The lowest order diagrams for $B_c \rightarrow J/\psi K$ decay.

where $r = M_{J/\psi}/M_{B_c}$, $r_b = M_b/M_{B_c}$, $r_c = M_c/M_{B_c}$ and

$$\begin{aligned} t_a^1 &= \max(M_B \sqrt{(1-x_2)(r_c-r^2)}, M_B \sqrt{|r_b^2 - (1-r^2)x_2|}, 1/b_1, 1/b_2), \\ t_a^2 &= \max(M_B \sqrt{r_c^2 + r_c - r^2}, 1/b_1, 1/b_2). \end{aligned} \quad (5)$$

The functions

$$\begin{aligned} h_a^{(1)}(x_2, b_1, b_2) &= S_t(x_2) K_0(M_B \sqrt{(1-x_2)(r_c-r^2)} b_1) \\ &\times [\theta(b_2 - b_1) \theta(r_b^2 - (1-r^2)x_2) I_0(M_B \sqrt{r_b^2 - (1-r^2)x_2} b_1) \\ &\times K_0(M_B \sqrt{r_b^2 - (1-r^2)x_2} b_2) + (b_1 \leftrightarrow b_2)], \end{aligned} \quad (6)$$

$$\begin{aligned} h_a^{(2)}(x_2, b_1, b_2) &= S_t(r_c) K_0(M_B \sqrt{(1-x_2)(r_c-r^2)} b_2) \\ &\times [\theta(b_2 - b_1) I_0(M_B \sqrt{r_c^2 + r_c - r^2} b_1) K_0(M_B \sqrt{r_c^2 + r_c - r^2} b_2) + (b_1 \leftrightarrow b_2)], \end{aligned} \quad (7)$$

come from the Fourier transformation of propagators of virtual quark and gluon in the hard part calculations. For the non-factorizable diagrams (c) and (d), all three meson wave functions are involved. Their total contribution is:

$$\begin{aligned} M_c[C] &= -\frac{16}{N_c} \pi C_F f_{B_c} M_{B_c}^2 \int_0^1 dx_2 dx_3 \int_0^\infty b_2 db_2 b_3 db_3 \phi_K^A(x_3) \\ &\times \{[(1-r^2)(1-r^2-r_c-x_3+r^2 r_c+r^2 x_3) \phi_\psi^L(x_2, b_2) + r(1-r^2)(1-r_c-x_2) \phi_\psi^t(x_2, b_2)] \\ &\alpha_s(t_c^1) h_c^{(1)}(x_2, x_3, b_2, b_3) \exp[-S_{B_c}(t_c^1) - S_{J/\psi}(t_c^1) - S_K(t_c^1)] C(t_c^1) + [(r^2-1)(1+r^2-2r_c-x_2+x_3 \\ &-r^2 x_2 - r^2 x_3) \phi_\psi^L(x_2, b_2) + r(1-r^2)(1-r_c-x_2) \phi_\psi^t(x_2, b_2)] \alpha_s(t_c^2) h_c^{(2)}(x_2, x_3, b_2, b_3) \\ &\exp[-S_{B_c}(t_c^2) - S_{J/\psi}(t_c^2) - S_K(t_c^2)] C(t_c^2)\}, \end{aligned} \quad (8)$$

where

$$\begin{aligned} t_c^1 &= \max(M_B \sqrt{|F_{(1)}^2|}, M_B \sqrt{(r_c-r^2)(1-x_2)}, 1/b_2, 1/b_3), \\ t_c^2 &= \max(M_B \sqrt{|F_{(2)}^2|}, M_B \sqrt{(r_c-r^2)(1-x_2)}, 1/b_2, 1/b_3), \end{aligned} \quad (9)$$

$$\begin{aligned} h_c^{(j)}(x_2, x_3, b_2, b_3) &= \\ &\left\{ \theta(b_2 - b_3) I_0(M_B \sqrt{(1-x_2)(r_c-r^2)} b_3) K_0(M_B \sqrt{(1-x_2)(r_c-r^2)} b_2) \right. \\ &\left. + (b_2 \leftrightarrow b_3) \right\} \times \begin{pmatrix} K_0(M_B F_{(j)} b_3), & \text{for } F_{(j)}^2 > 0 \\ \frac{\pi i}{2} H_0^{(1)}(M_B \sqrt{|F_{(j)}^2|} b_3), & \text{for } F_{(j)}^2 < 0 \end{pmatrix}, \end{aligned} \quad (10)$$

and $F_{(j)}$'s are defined by

$$\begin{aligned} F_{(1)}^2 &= (1 - r_c - x_3 + r^2 x_3)(x_2 - 1), \\ F_{(2)}^2 &= (1 - x_2)(r_c - r^2 - x_3 + r^2 x_3). \end{aligned} \quad (11)$$

The total decay amplitude is then

$$\mathcal{A}(B_c^+ \rightarrow J/\psi K^+) = V_{us} V_{cb}^* \left[C_1 M_c + \left(\frac{1}{3} C_1 + C_2 \right) M_a f_K \right], \quad (12)$$

and the decay width is expressed as

$$\Gamma(B_c^+ \rightarrow J/\psi K^+) = \frac{G_F^2 M_{B_c}^3}{128\pi} (1 - r^2) |\mathcal{A}(B_c^+ \rightarrow J/\psi K^+)|^2. \quad (13)$$

The following parameters have been used in our numerical calculation [11–15]:

$$\begin{aligned} M_{B_c} &= 6.286 \pm 0.005 \text{ GeV}, M_b = 5.2 \text{ GeV}, M_{J/\psi} = 3.097 \text{ GeV}, M_c = 1.82 \text{ GeV}, \\ f_{B_c} &= 489 \pm 4 \text{ MeV}, f_K = 160 \text{ MeV}, f_{J/\psi} = 405 \pm 14 \text{ MeV}, \\ \tau_{B_c} &= (0.46 \pm 0.07) \times 10^{-12} \text{ s}, |V_{us}| = 0.2257 \pm 0.0021, |V_{cb}| = (41.6 \pm 0.6) \times 10^{-3}. \end{aligned} \quad (14)$$

If not specified, we shall take their central values as the default input. We have taken the constituent quark masses $M_b = 5.2 \text{ GeV}$ and $M_c = 1.82 \text{ GeV}$ from the ISGW2 model[12]. As noted in Ref.[16], this choice for M_b and M_c satisfies approximately the relation $M_b = M_c + 3.4 \text{ GeV}$, which is consistent with the well known formula relating the pole masses $M_{b,pole}$ and $M_{c,pole}$ in the Heavy Quark Effective Theory. We also update some parameters used in previous works by taking the values from the latest Particle Data Group publication and lattice QCD simulations.

Our numerical analysis shows that $|C_1 M_c / ((\frac{1}{3} C_1 + C_2) M_a f_K)| = 17\%$, which means that the dominated contributions to the branching ratio of $B_c \rightarrow J/\psi K$ decays come from the factorizable topology[(a) and (b) in Fig.1]. We list our numerical results on the branching ratio for $B_c \rightarrow J/\psi K$ decays in Table I, where ω is the parameter in the wave function of the J/ψ meson(see Ref.[7] for details). From the numbers in Table I, one can find that the branching ratio of $B_c \rightarrow J/\psi K$ decays is sensitive to the parameter ω . The branching ratio of $B_c \rightarrow J/\psi K$ decays is more sensitive to the quark masses, especially the c-quark's mass, as shown in Table II. Therefore the $B_c \rightarrow J/\psi K$ decays provide a good platform to understand the wave function of the J/ψ meson and the constituent quark model. Besides the uncertainty from the parameter ω and the quark masses, we find the uncertainty of the decay constant $f_{J/\psi}$ will bring about 7% uncertainty to the branching ratio of $B_c \rightarrow J/\psi K$ decays. We investigate the branching ratio's dependence on the hard scale t in Eq. (3), which characterize the size of next-to-leading order contribution. The branching ratio is shown in Table III with those uncertainties. By changing the hard scale t from $0.8t$ to $1.2t$, we find the branching ratio for $B_c \rightarrow J/\psi K$ decays changes little as shown in Table III, that mean the uncertainties in the next-to-leading order contributions can be neglected for this decay mode. The value of Λ_{QCD} also affect the branching ratio of $B_c \rightarrow J/\psi K$ decays, we have taken $\Lambda_{QCD} = 250 \text{ MeV}$ at $N_f = 4$ as our default input. A recent determination of Λ_{QCD} gives $\Lambda_{QCD}^{N_f=4} = 234 \pm 26 \text{ MeV}$ [17]. Our result is $Br(B_c \rightarrow J/\psi K) = 1.70 \times 10^{-4}$, if $\Lambda_{QCD} = 208 \text{ MeV}$; and $Br(B_c \rightarrow J/\psi K) = 2.02 \times 10^{-4}$, if $\Lambda_{QCD} = 260 \text{ MeV}$. Considering the uncertainties from the input parameters, our results on the branching ratio of $B_c \rightarrow J/\psi K$ decays are generally in the range of $(1 - 3) \times 10^{-4}$. The large branching ratio and the clear signals of the final states make the measurement of $B_c \rightarrow J/\psi K$ easily at the LHC-b experiments.

	$\omega = 0.5 \text{ GeV}$	$\omega = 0.6 \text{ GeV}$	$\omega = 0.7 \text{ GeV}$
$Br(B_c^+ \rightarrow J/\psi K^+)$	1.52	1.96	2.44

TABLE I: Branch ratios in the unit 10^{-4} for different ω

In the literature, there already exist a lot of studies on $B_c \rightarrow J/\psi K$ decays[16, 18–24], we show their results in Table IV. All the previous works on this decay model were based on naive factorization, which, as expected, can be quite accurate for the B_c meson, since the quark-gluon sea is suppressed in the heavy quarkonium[20]. In the naive factorization approach, the decay amplitude can be expressed in terms of the hadronic transition form factors and of the leptonic decay constants. For the $B_c \rightarrow J/\psi$ transition form factors, there is a large difference among the previous works[7]. The authors of Ref.[19, 20] calculated the form factors in the frame work of QCD sum rules, as argued by

	$M_c = 1.72\text{GeV}$	$M_c = 1.82\text{GeV}$	$M_c = 1.92\text{GeV}$
$Br(B_c^+ \rightarrow J/\psi K^+)$	3.80	1.96	1.19

TABLE II: Branch ratios in the unit 10^{-4} for different M_c using $\omega = 0.6\text{GeV}$ and $M_b = 5.2\text{GeV}$

Scale	$Br(B_c^+ \rightarrow J/\psi K^+)$
$0.8\ t$	2.10×10^{-4}
t	1.96×10^{-4}
$1.2\ t$	1.91×10^{-4}

TABLE III: Branching ratio using $\omega = 0.6\text{GeV}$ with uncertainties

the authors, the sum rule estimates of form factors are taken at zero transfer squared, while the dependence on q^2 is beyond the reliable accuracy of the method. The authors of Ref.[16, 22–24] presented their calculations based on the constituent quark model, where, the B_c decay form factors can be expressed through the overlap integrals of the meson wave functions. In Ref.[21], Chang and Chen used a very complicated approach to evaluate the weak current matrix elements. It is expected that the approach is available as long as the mesons in the initial and final states are of weak binding, for example, for the decay $B_c \rightarrow J/\psi K$. After updating the model parameters, Ivanov *et al.* studied exclusive nonleptonic and semileptonic decays of the B_c meson within a relativistic constituent quark model in Ref.[18]. For the branching ratio of $B_c \rightarrow J/\psi K$ decays, their results are close to our predictions. In general, the results of the various model calculations on $B_c \rightarrow J/\psi$ decays are of the same order of magnitude. Eventually, the variations among the theoretical predictions should be discerned by experimental results of the B_c decays, where, one can learn much about the decay mechanism.

This work is supported by the National Natural Science Foundation of China under Grant Nos. 10847157 and 10575083. We would to thank Dr. M.-Z. Zhou and Dr. J.-F. Sun for valuable discussions.

-
- [1] N. Brambilla, *et al.*, (Quarkonium Working Group), CERN-2005-005, hep-ph/0412158.
 - [2] M. P. Altarelli and F. Teubert, Int. J. Mod. Phys. **A23**, 5117(2008).
 - [3] H.-n. Li and H. L. Yu, Phys. Rev. Lett. **74**, 4388 (1995); Phys. Lett. **B353**, 301 (1995);
H.-n. Li, *ibid.* 348, 597 (1995); H. n. Li and H. L. Yu, Phys. Rev. **D53**, 2480 (1996).
 - [4] G. Buchalla, A. J. Buras and M. E. Lautenbacher, Rev. Mod. Phys. **68**, 1125(1996).
 - [5] C.-D. Lü, K. Ukai, and M.-Z. Yang, Phys. Rev. **D63**, 074009(2001).
 - [6] J. Zhang and X.-Q. Yu, Eur. Phys. J. **C63**, 435(2009).
 - [7] Jun-Feng Sun, Dong-Sheng Du and Yue-Ling Yang, Eur. Phys. J. **C60**, 107(2009).
 - [8] J.-F. Cheng, D.-S. Du and C.-D. Lü, Eur. Phys. J. **C45**, 711(2006).
 - [9] X.-Q. Yu, Y. Li and C.-D. Lü, Phys. Rev. **D71**, 074026(2005).
 - [10] A. Khodjamirian, T. Mannel and M. Melcher, Phys. Rev. **D70**, 094002 (2004); V.M. Braun and A. Lenz, Phys. Rev. **D70**, 074020 (2004).
 - [11] C. Amsler, *et al.*(Particle Data Group), Phys. Lett. **B667**, 1(2008).
 - [12] D. Scora and N. Isgur, Phys. Rev. **D52**, 2783(1995).
 - [13] T. W. Chiu, T. H. Hsieh(TWQCD Collaboration), PoS. **LAT2006**, 180(2007).
 - [14] B. Melic, Phys. Rev. **D68**, 034004(2003).
 - [15] S. Hashimoto, Int. J. Mod. Phys. **A20**, 5133(2005).
 - [16] A. Y. Anisimov, I. M. Narodetskii, C. Semay and B. Silvestre-Bra, Phys. Lett. **B452**, 129(1999); A. Y. Anisimov, P. Y. Kulikov, I. M. Narodetskii and K. A. Ter-Martirosyan, Phys. Atom. Nucl. **62**, 1739(1999).
 - [17] J. Blumlein, arXiv:0706.2430.

	[18]	[19, 20]	[21]	[16, 22]	[23]	[24]
$Br(B_c \rightarrow J/\psi K)$	1.3×10^{-4}	1.1×10^{-4}	1.4×10^{-4}	0.7×10^{-4}	0.5×10^{-4}	0.8×10^{-4}

TABLE IV: Branching ratio for $B_c \rightarrow J/\psi K$ decay in the literature.

- [18] M. A. Ivanov, J. G. Korner and P. Santorelli, Phys. Rev. **D73**, 054024(2006).
- [19] V. V. Kiselev, A. E. Kovalsky and A. K. Likehoded, Nucl. Phys. **B585**, 353(2000); arXiv:hep-ph/0006104.
- [20] V. V. Kiselev, arXiv:hep-ph/0211021.
- [21] C. H. Chang and Y. Q. Chen, Phys. Rev. **D49**, 3399(1994).
- [22] P. Colangelo and F. De Fazio, Phys. Rev. **D61**, 034012(2000).
- [23] D. Ebert, R. N. Faustov and V. O. Galkin, Mod. Phys. Lett. **A17**, 803(2002); Eur. Phys. J. **C32**, 29(2003); Phys. Rev. **D68**, 094020(2003).
- [24] A. Abd El-Hady, J. H. Munoz and J. P. Vary, Phys. Rev. **D62**, 014019(2000).